

# Comprehensive Cardiovascular Assessment

sample type: **BLOOD**

**Comprehensive Cardiovascular Assessment** analyses blood for lipid markers, ratios, and independent risk factors. Together, these markers provide a thorough assessment of cardiovascular health status, revealing the biochemical environment associated with inflammation, lipid deposits, endothelial dysfunction, clotting factors and other primary mechanisms underlying cardiovascular disease.

Nearly 50% of all heart attack victims have normal levels of traditional early warning markers for cardiovascular disease (CVD), including total cholesterol. For this reason, improved clinical tools are needed to identify patients with a predisposition to CVD who can benefit from timely, preventative intervention. The **Comprehensive Cardiovascular Assessment** measures state-of-the-art, independent markers that illuminate the clinical complexity of CVD.

## Markers for CVD

- **Homocysteine** is an amino acid that functions as an intermediate in methionine metabolism. It can rise in response to nutritional deficiencies of B12, folate, B6, or betaine. High levels of homocysteine have been linked to damaged endothelium, increased platelet aggregation, and the formation of atherosclerotic lesions.
- **Lipoprotein(a) or Lp(a)** is a marker strongly influenced by heredity. It has been cited for its causative role in atherothrombogenesis and its strong association with both coronary and peripheral cardiac events. Lp(a) promotes the deposit of fatty wastes and other debris in atherosclerotic lesions.
- **C-reactive protein (CRP)** is an important independent marker for inflammation. High levels reflect overactivity of inflammatory cytokines linked to coagulation and vascular endothelium damage. Evidence suggests that previous infection with pathogens such as Chlamydia pneumoniae or Helicobacter pylori may act as an initiating trigger for this chronic inflammation. Elevated CRP has also been linked to a poorer long-term prognosis in individuals with a recent history of a cardiac event.
- **Fibrinogen** plays a key role in arterial occlusion by promoting thrombus formation, endothelial injury and hyperviscosity.
- **Blood Lipids** - Triglycerides may become elevated by diets containing excess amounts of saturated fats or carbohydrates. These elevations are implicated in the progression of both coronary and peripheral atherosclerosis. HDL and LDL cholesterol, Apo-A1, Apo-B, total cholesterol and their ratios assess specific dynamics of lipid metabolism and their potential impact on CVD.
- **Glycosylated haemoglobin** is a measure of red blood cells damaged by glucose, reflecting long-term imbalanced blood-sugar levels. This is especially important in those with metabolic syndrome or diabetes where the risk of cardiovascular disease is significantly increased.

A composite **Cardiovascular Index** provides an overall indication of synergistic risk. The index incorporates a multivariate analysis of all the individual lipid scores and the independent CV factors.

A wide range of nutritional, lifestyle, and dietary interventions can be used to improve the patient's biochemical profile and optimise cardiovascular health.

- **Analytes:**  
lipoprotein(a)  
homocysteine  
fibrinogen  
C-reactive protein  
glycosylated haemoglobin  
triglycerides  
total cholesterol  
LDL cholesterol  
HDL cholesterol  
apolipoprotein B  
apolipoprotein A-1  
total cholesterol/HDL  
Triglycerides/HDL  
Apo B/Apo A-1  
cardiovascular index

- **Specimen Requirements:**  
- 3 ml plasma and  
- 4 ml serum, fasting

- **Before Taking this Test:**  
- Fast overnight (at least 12 hours)  
- Inform practitioner about medication and supplement use, including aspirin and cholesterol-lowering drugs  
- See instructions inside test kit for details

- \* **NOTE!**—Samples must be processed & centrifuged within 15 minutes, therefore please arrange with referring phlebotomist or arrange collection at our phlebotomy locations in Surrey or London  
-- **Arrange blood draw Monday to Wednesday only**



Genova  
Diagnostics®  
Europe

Innovative Testing for Optimal Health

Accession No:



# Comprehensive Cardiovascular Assessment

## Patient Details

Ms Sample Report  
 Parkgate House  
 356 West Barnes Lane  
 New Malden  
 Surrey  
 KT3 6NB

## Practitioner Details

Genova Diagnostics ( Europe )  
 Parkgate House  
 356 West Barnes Lane  
 New Malden  
 Surrey  
 KT3 6NB

Client ID No: IWX500220  
 Accession No:  
 Patients DOB: 20/03/1975  
 Sample Date:  
 Date Of Report: 06/02/2009

This test reveals important clinical information about:


- **Standard lipid risk markers for cardiovascular disease (CVD) risk**, including triglycerides, total cholesterol, LDL cholesterol, and Apo B
- **Protective factors HDL cholesterol and Apo A-1**, along with significant lipid ratios
- **Independent risk factors for CVD risk**, including C-reactive protein, homocysteine, Lp(a), and fibrinogen
- **Synergy of risk factors**, reflected by an overall Cardiovascular Risk Index

Lipid Markers		
Analyte	Result	Reference Range
Triglycerides	0.55	<= 1.69 mmol/L
Total Cholesterol	5.20 H	3.9 - 5.17 mmol/L
LDL Cholesterol	3.00 H	<= 2.59 mmol/L
HDL Cholesterol	1.60	>= 1.55 mmol/L

Independent CV Factors		
Analyte	Result	Reference Range
Hb A1c	4.50	4.3 - 6.1 % of Hb
Homocysteine	4.34	3 - 10 µmol/L
hs-CRP	1.22 H	<= 1.00 mg/L
Fibrinogen	8.76	5.3- 10.3 µmol/L

Lipoprotein Markers		
Analyte	Result	Reference Range
Apo A-1	1.68	1.25 - 2.15 g/L
Apo B	0.91	0.55 - 1.25 g/L
Lp (a)	0.39	<= 1.14 µmol/L

Ratios		
Analyte	Result	Reference Range
Total Cholesterol / HDL	3.25	<= 3.7
Triglycerides / HDL	0.79	<= 3.0
Apo B / Apo A-1	0.54	0.30 - 0.90



For test kits, clinical support, or more information contact:

Genova Diagnostics, Europe  
 Parkgate House  
 356 West Barnes Lane  
 New Malden  
 Surrey  
 KT3 6NB

+44 (0)20 8336 7750 • Fax: +44 (0)20 8336 7751

More detailed publications with references are also available: [www.GDXuk.net](http://www.GDXuk.net)